

This leaflet has been produced for your information.  
For further advice please contact the school office or  
your doctor's surgery.

**Chickenpox is common and mostly affects children, although you can get it at any age. It usually gets better by itself within a week without needing to see a GP.**

### **Symptoms of Chickenpox**

1. Chickenpox starts with red spots. They can appear anywhere on the body.
2. The spots fill with fluid and become blisters. The blisters may burst. They might spread or stay in a small area.
3. The spots scab over. More blisters might appear while others scab over.

### **Other symptoms**

You might get symptoms before or after the spots, including:

- a high temperature above 38C
- aches and pains, and generally feeling unwell
- loss of appetite

Chickenpox is very itchy and can make children feel miserable, even if they do not have many spots. Chickenpox is usually much worse in adults.

It's possible to get chickenpox more than once, although it's unusual.



## **Important**



**You'll need to stay away until all the spots have crusted over. This is usually 5 days after the spots first appeared.**

## **Do**

- drink plenty of fluid (try ice lollies if your child is not drinking) to avoid dehydration
- take [paracetamol](#) to help with pain and discomfort
- put socks on your child's hands at night to stop scratching
- cut your child's nails
- use cooling creams or gels from your pharmacy
- speak to a pharmacist about using antihistamine medicine to help itching
- bathe in cool water and pat the skin dry (do not rub)
- dress in loose clothes

## **Don't**

- do not use [ibuprofen](#) unless advised to do so by your doctor, as it may cause serious skin infections
- do not give aspirin to children under 16
- do not be around pregnant women, newborn babies and people with a weakened immune system, as it can be dangerous for them

## **Non-urgent advice: Speak to a GP if:**

- you're not sure it's chickenpox
- the skin around the blisters is red, hot or painful (signs of infection)
- your child is [dehydrated](#)
- you're concerned about your child or they get worse

Tell the receptionist you think it's chickenpox before going in. They may recommend a special appointment time if other patients are at risk.

### **It's easy to catch chickenpox**

You can catch chickenpox by being in the same room as someone with it.

It's also spread by touching clothes or bedding that has fluid from the blisters on it.

### **How long chickenpox is infectious for**

Chickenpox is infectious from 2 days before the spots appear to until they have crusted over, usually 5 days after they first appeared.

### **How soon you get symptoms after coming into contact with chickenpox**

It takes 1 to 3 weeks from the time you were exposed to chickenpox for the spots to start appearing